



REVIEW

Mechanisms of non-pharmacologic adjunct therapies used during exercise in COPD

A.M. Moga^{a,b}, M. de Marchie^c, D. Saey^d, J. Spahija^{a,b,e,*}

^a School of Physical and Occupational Therapy, McGill University, 3654 Promenade Sir William Osler, Montreal, Quebec H3G 1Y5, Canada

^b Respiratory Health Research Center, Sacré-Coeur Hospital, 5400 boul. Gouin Ouest, Montréal, , Quebec H4J 1C5, Canada

^c Department of Adult Critical Care, Sir Mortimer B. Davis Jewish General Hospital, McGill University, Montreal, Quebec, Canada

^d Centre de Recherche, Institut Universitaire de Cardiologie et de Pneumologie de Québec, Université Laval, Québec, Canada

^e Center for Interdisciplinary Research in Rehabilitation in Montreal, Jewish Rehabilitation Hospital, 3205, Place Alton-Goldbloom, Laval, Quebec H7V 1J1, Canada

Received 8 August 2011; accepted 12 January 2012
Available online 15 February 2012

KEYWORDS

COPD;
Exercise;
Non-invasive
mechanical
ventilation;
Heliox;
Supplemental
oxygen;
Respiratory
mechanics

Summary

Individuals with chronic obstructive pulmonary disease (COPD) are often limited in their ability to perform exercise due to a heightened sense of dyspnea and/or the occurrence of leg fatigue associated with a reduced ventilatory capacity and peripheral skeletal muscle dysfunction, respectively. Pulmonary rehabilitation programs have been shown to improve exercise tolerance and health related quality of life. Additional therapeutic approaches such as non-invasive ventilatory support (NIVS), heliox (He–O₂) and supplemental oxygen have been used as non-pharmacologic adjuncts to exercise to enhance the ability of patients with COPD to exercise at a higher exercise-intensity and thus improve the physiological benefits of exercise. The purpose of the current review is to examine the pathophysiology of exercise limitation in COPD and to explore the physiological mechanisms underlying the effect of the adjunct therapies on exercise in patients with COPD. This review indicates that strategies that aim to unload the respiratory muscles and enhance oxygen saturation during exercise alleviate exercise limiting factors and improve exercise performance in patients with COPD. However, available data shows

* Corresponding author. Hôpital du Sacré-Cœur de Montréal, Axe de recherche en santé respiratoire, 5400 boul. Gouin Ouest, Montréal, Quebec H4J 1C5, Canada. Tel.: +1 5143382222x3654; fax: +1 5147397357.

E-mail address: jadranka.spahija@mcgill.ca (J. Spahija).

significant variability in the effectiveness across patients. Further research is needed to identify the most appropriate candidates for these forms of therapies.

© 2012 Elsevier Ltd. All rights reserved.

Contents

Introduction	615
Exercise limitation in COPD	616
Ventilatory limitation and work of breathing	616
Peripheral muscle dysfunction	617
Cardiac function and blood distribution	618
Non-pharmacologic adjunct therapies	618
Non-invasive ventilatory support	618
Effect of NIVS on respiratory and peripheral muscles and their interaction	619
Heliox	619
Effect of heliox on respiratory and peripheral muscles and their interaction	620
Supplemental oxygen	620
Effect of supplemental oxygen on the ventilatory and peripheral muscles and their interaction	620
Conclusion	621
Conflict of interest	621
References	621

Introduction

Chronic obstructive pulmonary disease (COPD) is a pulmonary disorder that is characterized by progressive irreversible airflow limitation resulting from alveolar wall destruction, bronchiolar narrowing¹ and airway inflammation that occurs in response to inhalation of noxious particles or gases.² Although numerous genetic, occupational, and environmental factors have been associated with COPD,^{3–5} cigarette smoking remains the primary cause of the disease.⁶ COPD is a major cause of morbidity and mortality and poses a substantial economic and social burden worldwide.^{1,2}

Individuals with COPD commonly exhibit a limited ability to perform exercise.^{2,7,8} Compared to healthy individuals, patients with COPD demonstrate lower maximum exercise capacities and lower levels of peak oxygen consumption ($\text{VO}_{2\text{peak}}$),^{9–11} with the lowest levels observed in patients with more severe COPD.^{10–12} Although moderate correlations between $\text{VO}_{2\text{peak}}$ and the force expiratory volume in the first second (FEV_1) have been reported in patients with mild ($r = 0.69$), moderate ($r = 0.65$), and severe ($r = 0.87$) COPD, others have found FEV_1 to be a poor predictor of exercise capacity.^{13–15} Patients with COPD typically experience dyspnea during exercise; however, the locus of symptom limitation (i.e., the reason for stopping exercise) is not uniform across patients.¹⁶ Whereas the majority of patients with COPD stop exercise because of dyspnea, others are limited by leg fatigue or a combination of dyspnea and leg fatigue. Compared to individuals with mild COPD, those with moderate-to-severe disease tend to perceive dyspnea more intensely than leg fatigue.¹⁷ However, patients with COPD also exhibit skeletal muscle

abnormalities, which can contribute to exercise intolerance.¹⁸ Although the exact proportion varies among studies, leg fatigue has been reported as the primary symptom limiting exercise during cycling in approximately one third of the patients with COPD.^{17,19} Some studies have also reported a moderate correlation between leg discomfort during exercise and the magnitude of contractile muscle fatigue in patients with COPD.^{19,20}

In an effort to help reduce dyspnea and improve exercise capacity, individuals with COPD are often referred to pulmonary rehabilitation programs.^{21–24} Such programs typically use a multidisciplinary approach, combining education and exercise to optimize physical and social performance and autonomy. However, exercise training has been shown to be the essential component for improving exercise capacity and health related quality of life (HRQoL).^{21–23,25–28} The physiological benefits associated with high intensity exercise training include a reduction of exercise lactic acidosis and heart rate for a given work rate,²⁹ which in turn leads to a lower ventilatory demand and a more effective breathing pattern,³⁰ enhanced activity of mitochondrial enzymes and capillary density in the trained muscles,^{30,31} as well as enhanced anabolic processes in the peripheral muscles.³² There is also some evidence that whole body aerobic exercise training may improve respiratory muscle function in patients with COPD, as demonstrated by an increase in the maximum inspiratory muscles pressure.^{33–35} While low intensity exercise training has also been shown to be effective in improving exercise tolerance with regard to endurance for activities such as walking, it does not lead to the same physiologic training effect that can result from high intensity training.²⁹ Although studies suggest that

most patients with COPD can benefit from pulmonary rehabilitation,^{2,27,29,30,36,37} some individuals with severe lung disease may be unable to obtain a true physiological training effect because of their inability to exercise at a high enough exercise-intensity (i.e., 80% of maximum).^{2,29}

Therapeutic approaches such as non-invasive ventilatory support (NIVS),^{22,38,39} low-density gases (i.e. Heliox),^{40,41} and supplemental oxygen (O₂)^{42,43} have been used as non-pharmacologic adjuncts to exercise training to enhance the ability of patients with COPD to exercise at a higher exercise-intensity and thus improve the physiological benefits of exercise.

The current review examines the factors that contribute to exercise limitation in COPD and the physiological mechanisms via which non-pharmacologic adjunct therapies may improve exercise tolerance. Understanding how these adjunct therapies affect the factors contributing to exercise limitation is of paramount importance for guiding identification of the most appropriate forms of therapy for improving exercise in such individuals.

Exercise limitation in COPD

Ventilatory limitation and work of breathing

The changes that occur to the mechanical properties of the lungs in COPD contribute to an increased work of breathing^{44,45} and impaired gas exchange.² Expiratory flow limitation resulting from airway inflammation and loss of lung elasticity⁴⁶ leads to air trapping within the lungs which increases the end-expiratory lung volume (i.e. static pulmonary hyperinflation).^{47–49} During exercise, minute ventilation (V_E) is increased predominantly by an increased respiratory rate (RR), whereas tidal volume (V_T), which approaches the limits of total lung capacity at end-inspiration, increases marginally before reaching a plateau (Fig. 1).⁵⁰ The increased RR results in less time available for exhalation, and leads to further increases in end-expiratory lung volume (i.e. dynamic hyperinflation).^{49–52} Because the respiratory system is less compliant at higher lung volumes, static and dynamic hyperinflation contribute to an increase in the inspiratory elastic work of breathing. This is further compounded by an




Figure 1 Ventilatory response to incremental exercise is shown in patients with COPD and in age-matched healthy individuals. Changes in dyspnea intensity (upper left), operational lung volumes (upper right), breathing pattern (lower left) and the effort-displacement ratio (lower right) are shown as ventilation increases with exercise. Dyspnea intensity is greater and breathing pattern is more rapid and shallow in COPD (solid circles) compared to healthy individuals (open circles). Greater mechanical constraints on tidal volume (VT) are evident in COPD because of increasing end-expiratory lung volumes and limitation from above by the total lung capacity (TLC). To increase ventilation, individuals with COPD increase breathing frequency (F) to a greater extent. Tidal inspiratory pressure swings expressed as a fraction of their maximal force-generating capacity ($P_{es}/P_{I_{max}}$) relative to the VT response expressed as a fraction of the predicted vital capacity (VC) show a significantly steeper slope in COPD.⁸ Reprinted with permission of the American Thoracic Society. Copyright (©) 2012 American Thoracic Society. Official Journal of the American Thoracic Society.

inspiratory threshold load whereby patients with COPD must often generate an inspiratory threshold pressure before airflow occurs, i.e. intrinsic positive end-expiratory pressure (PEEPi).⁵³

The development of hyperinflation additionally decreases inspiratory muscle length, which reduces the ability of such muscles to generate force.⁵⁴ Dynamic hyperinflation therefore contributes to neuromechanical uncoupling whereby an increased inspiratory effort is required to generate a given ventilatory output. Under conditions of impaired length–tension relationship⁵⁴ and reduced coupling, COPD patients need to increase central respiratory drive and diaphragm activation^{55–57} in order to maintain the same pressure generated across the diaphragm (i.e. transdiaphragmatic pressure)⁵⁷ (Fig. 2). This increased activation has been associated with greater respiratory effort sensation,^{48,50,56} increased energy demands,^{58,59} as well as enhanced inspiratory muscle fatigability.^{57,58,60,61}

Peripheral muscle dysfunction

In addition to the clear evidence of dynamic hyperinflation and altered pulmonary mechanics contributing to exercise intolerance in COPD, peripheral muscle dysfunction also appears to play an important role in limiting exercise in such individuals.^{2,17,62}

The skeletal muscle abnormalities that occur in COPD result from both deconditioning⁶³ secondary to decreased activity levels,³¹ and from systemic inflammation.^{64,65} These abnormalities are characterized by a reduction in

the proportion of type I fatigue-resistant fibers,^{66,67} increased proportion of less efficient type II fibers,^{68,69} reduced cross-sectional area of type I fibers i.e. muscle atrophy,^{66,68} and a reduced oxidative enzyme activity.⁷⁰ Several studies have shown a relationship of such changes with reduced peripheral muscle strength and endurance, as well as an increased contractile fatigability.^{62,68,70,71} Contractile muscle fatigue has been defined as a reversible reduction in the capacity of the skeletal muscle to generate force in response to a given neural input.^{72,73} Evidence for the occurrence of quadriceps contractile fatigue during exercise in patients with COPD comes from several studies that have assessed quadriceps muscle strength via magnetic stimulation.^{74–78} Studies have shown quadriceps strength to be correlated with maximum exercise capacity, independent of pulmonary function.^{71,77,79} Saey et al.⁷⁷ demonstrated that even after an improvement in FEV₁ following bronchodilator medication, quadriceps contractile fatigue was still evident following endurance cycling in a subgroup of patients who previously reported leg fatigue as being the primary factor limiting maximum exercise. These findings suggest that the presence of leg fatigue modulates the exercise response to bronchodilation.

Although muscle fatigue is a complex phenomenon, changes in muscle energy metabolism may be involved.⁷⁷ Several studies have found that compared with age-matched healthy individuals, COPD patients have lower lactate thresholds (i.e. VO₂ at which blood lactic acid begins to increase).^{63,80} The lower lactate thresholds lead to an excessive accumulation of metabolic by-products during exercise,^{29,81} which further impairs contractility




Figure 2 Campbell diagram showing the volume of lung or chest wall (V_L) plotted against pleural pressure (P_{Pi}) and the effects of dynamic hyperinflation on inspiratory muscle work. FRC: functional residual capacity. The continuous line (loop) traces a complete breath from FRC; arrows show direction. The dashed line shows the elastic characteristic of the lung (negative of elastic recoil pressure), and the dotted line shows the elastic characteristic of the relaxed chest wall. P_{mus} , pressure change generated by inspiratory muscles (length of the horizontal arrow). PEEPi, intrinsic positive end-expiratory pressure that must be overcome before inspiratory flow can begin (length of the horizontal arrow). The diagonally hatched area is work done against resistance, stippled area is work done against elastance of lung and chest wall, and horizontally hatched area is work to overcome PEEPi. (A) Illustrates the FRC is equal to V_{rel} , the relaxation volume of the respiratory system in a healthy individual, and (B) FRC is increased above relaxation volume in an individual with COPD. In COPD, the work done against inspiratory resistance is increased whereas the elastic characteristics of lung and chest wall are unchanged. Ref. 53.

and increases the risk of fatigue in patients with COPD.¹⁸ The increased CO₂ production also further increases the ventilatory demand and induces ventilatory limitation at lower than normal exercise workloads,⁸² thus causing early exercise termination in such patients.^{2,7,17}

Cardiac function and blood distribution

Cardiac output (CO), which is the body's energy supply and the product of stroke volume and heart rate (HR) is regulated principally by the demand for O₂ by the cells of the body.⁸³ In general, the exercise-induced increase in VO₂ is achieved by an increased CO and an increased O₂ extraction at the level of the working respiratory and peripheral skeletal muscles. During both sub-maximal and maximal exercise in healthy individuals, CO increases nearly linearly with VO₂, suggesting that O₂ consumption is linearly related to the energy supply.^{84–86} Although COPD patients likewise present an almost linear CO and VO₂ relationship during sub-maximal exercise,^{87–89} HR has been reported to be higher than normal at any given VO₂,⁹⁰ implying that stroke volume (SV) is lower than normal.⁹¹ In the absence of coexisting left-sided heart disease, evidence suggests that the decreased SV occurs secondary to reduced right ventricular ejection fraction at rest and which, on average, fails to increase with exercise among patients with COPD.^{92–94} As with CO, blood flow to the exercising peripheral muscles at a given sub-maximal exercise work rate is similar in patients with COPD and healthy individuals.^{11,81} In contrast, individuals with COPD reach lower peak exercise work rates and exhibit a reduced peak VO₂ (30–50% lower) and peak CO (35–60% lower) with a normal or reduced HR, as well as a lower peak leg blood flow compared to healthy individuals.^{9–11,87–89,95–97}

Dynamic hyperinflation appears to have a significant impact on cardiovascular function during exercise in patients with COPD.⁹¹ Vassaux et al.⁹⁸ showed a strong association ($r = 0.87$) between inspiratory-to-total lung capacity ratio (IC/TLC represents an index of hyperinflation) at rest and exercise and O₂ pulse (i.e. VO₂/HR, a surrogate marker of cardiac function), demonstrating that the most hyperinflated patients (i.e. IC/TLC $\leq 0.25\%$, an indicator of severe static hyperinflation), had a lower peak exercise O₂ pulse at a similar work load than patients having less hyperinflation (i.e., IC/TLC $> 0.25\%$).⁹⁸ In an attempt to maintain the required ventilation during exercise when the ability to increase O₂ supply is limited, individuals with severe hyperinflation are forced to generate a larger intra-thoracic pressure.^{92,99} This in turn, limits venous return, right and left ventricular (LV) blood volumes, and consequently, cardiac output.^{92,100,101} In addition, loss of pulmonary vascular capacity with emphysema results in an increased pulmonary vascular resistance which may ultimately impair LV filling. Barr et al.¹⁰¹ reported that the severity of airflow obstruction (FEV₁/FVC) and the degree of emphysema on chest CT scans was inversely correlated with reductions in LV end-diastolic volume, stroke volume and CO. Although there was a stronger association in more severe patients, hemodynamic changes also occurred with mild emphysema and airflow obstruction. Watz et al.¹⁰² reported that IC/TLC was

more strongly correlated with cardiac chamber size than measurements of airway obstruction or diffusion capacity, and that COPD patients with IC/TLC $\leq 0.25\%$ not only have an impaired LV filling, but also a reduced functional capacity as indicated by a lower 6-min walk distance.

Several studies have suggested^{10,103–105} that when the energy demands of the respiratory muscles are increased, such as during exercise, a competition for blood flow develops between the respiratory and peripheral muscles, which ultimately favors a redistribution of blood flow from the locomotor to the respiratory muscles. Evidence for this phenomenon, known as the "respiratory steal" or "blood stealing effect",¹⁰³ comes from Simon et al.,¹⁰⁶ who found that about 45% of the patients with COPD participating in their study demonstrated a leg blood flow plateau during whole body incremental cycling exercise, despite increasing exercise work load. They additionally found that the patients who exhibited such a leg blood flow plateau also revealed a greater work of breathing at sub-maximal exercise, indicating a high O₂ demand of the respiratory muscles.¹⁰⁶ In this context, it has been suggested that reduction in blood flow to the working peripheral muscles may induce leg fatigue, thereby limiting the duration and the intensity of exercise in patients with COPD as demonstrated by other studies.^{82,107}

Non-pharmacologic adjunct therapies

In the last several years, adjunct therapies to exercise training such as non-invasive ventilatory support and heliox have been investigated in an attempt to counter the high respiratory muscle workloads experienced by patients with COPD during exercise. It is anticipated that by unloading the inspiratory muscles, such strategies might enable individuals to exercise at higher intensities, thereby increasing also the training load to the peripheral muscles and enhancing the physiologic benefits of exercise training.

Non-invasive ventilatory support

Although non-invasive mechanical ventilation has traditionally been used with patients who have respiratory failure or sleep apnea,^{108,109} it has recently gained more attention as a potential tool for increasing exercise tolerance during pulmonary rehabilitation in patients with COPD.¹¹⁰ Non-invasive ventilatory support (NIVS) differs from traditional invasive mechanical ventilation, by the fact that it does not require the patient to be intubated (i.e. endotracheal or nasotracheal tube) for delivery of the positive pressure.¹¹¹ NIVS can be delivered through a variety of interfaces such as a mouthpiece, nasal prongs, or facemask.^{39,112,113}

The modes of mechanical ventilation that have been used for the delivery of NIVS during exercise include: (1) continuous positive airway pressure (CPAP), which delivers a constant positive pressure that elevates the baseline pressure (airway pressure which is constantly higher than atmospheric pressure)^{38,114}; (2) bilevel positive airway pressure (BiPAP) provides continuous positive pressure at two levels, a higher one for inspiration (IPAP) and a lower for expiration (EPAP), where both are above atmospheric

pressure, and the difference between IPAP and EPAP is a reflection of the amount of pressure support provided to the patient^{115–122}; (3) pressure support ventilation (PSV),^{123–127} which is a pressure-targeted mode whereby each breath is patient triggered and cycled; and (4) proportional assist ventilation (PAV)^{128–132} which provides assist in proportion to the patient's spontaneous effort, according to the equation of motion (requiring the determination of elastance and resistance and instantaneous measures of flow and volume). This requires that the pressure that is delivered within a breath is continuously readjusted in proportion to the pressure that is generated by the inspiratory muscles, determined using instantaneous measurements of inspiratory airflow and volume.¹²⁸

Effect of NIVS on respiratory and peripheral muscles and their interaction

There is evidence that NIVS unloads the inspiratory muscles and reduces the work of breathing both at rest and during exercise.^{127,133,134} NIVS has also been shown to decrease dyspnea, and increase endurance time in individuals with moderate-to-severe COPD.^{38,103,127,129,135,136} Previous studies have demonstrated a relationship between decreased dyspnea and reduced work of breathing^{127,135,136} as well as decreased dyspnea and diaphragm deactivation.¹³⁷ Unloading the respiratory muscles during high-intensity exercise (70–80%Wmax) using NIVS has also been found to improve peripheral muscle oxygenation^{117,138} and to reduce blood lactate levels^{129,139} which in turn not only reduces the occurrence of leg fatigue, but also further decreases respiratory drive and dyspnea,^{129,139} and thereby has the potential to decrease ventilatory limitation to exercise.

Several studies have demonstrated that NIVS administration during exercise increases V_E as a result of both increased V_T and RR^{127,129} or only V_T ,³⁸ whereas others have reported no change in V_E for a given work load.^{103,125,135,140} There is evidence, however, that NIVS promotes a reduction in inspiratory work load, whether¹²⁷ or not¹³⁵ V_T and end-inspiratory lung volume are increased.^{133,141} Although NIVS has no direct effect on end-expiratory lung volume,^{38,133,141} PSV during exercise has been reported to promote greater diaphragm muscle deactivation in patients with COPD compared to healthy individuals,¹³³ supporting the unloading effect of NIVS.

In patients with more severe COPD, up to 50% of the whole body VO_2 during exercise goes to the respiratory muscles due to an increased work of breathing,¹⁴² enhancing the likelihood of the occurrence of the respiratory steal phenomenon.^{10,103} Several lines of evidence suggest that reducing the work of breathing via NIVS may decrease ventilatory muscle blood flow requirements and allow a fraction of the limited CO to be redirected to the locomotor muscles, thereby improving peripheral muscle perfusion, and in turn exercise capacity.^{103,143} These responses to NIVS have been found to improve endurance time in both healthy individuals^{104,105} and patients with COPD.^{117,138}

The effects of NIVS on the cardiac performance are complex, with most resulting from an NIVS associated rise

in mean intra-thoracic pressure and a fall in transmural pressure.^{134,144–146} In a study evaluating the effect of PSV+PEEP, Oliveira et al.¹³⁸ showed that NIVS promoted an increase in stroke volume, HR, CO, and ultimately exercise endurance in one subgroup of COPD patients; however, in another subgroup, NIVS resulted in a decreased stroke volume and HR, thereby reduced CO with no improvement in exercise endurance. The study found that patients in the latter subgroup tended to be more hyperinflated, suggesting that NIVS may have a deleterious effect on hemodynamics and exercise tolerance in COPD patients who exhibit greater static hyperinflation.

Although there is no evidence to date supporting the use of one mode of NIVS over another, PAV has been advocated for exercise training because it is believed to enhance the synchrony between patient effort and ventilatory support^{128,147} thus improving patient comfort, reducing dyspnea and increasing exercise tolerance.^{129,130} However, the need for continuous measurement of the patient's respiratory mechanics (i.e., resistance, elastance, and iPEEP) and adjustment of ventilator settings greatly increases the complexity of this mode of mechanical ventilation.

From the accumulated evidence, there is ample empirical data showing that NIVS applied during exercise unloads the inspiratory muscles, decreases the drive to breathe thereby reducing dyspnea, delays lactate buildup, and ultimately improves exercise performance among patients with COPD. Such improvements, however, vary considerably among individuals. Moreover, there is inconsistent data corroborating the use of NIVS during routine pulmonary rehabilitation programs for increasing the overall benefit of pulmonary rehabilitation compared to training alone.¹³² Discomfort to ventilator settings and/or the interface used to deliver the assist¹²⁰ are factors that may contribute to lack of tolerance to NIVS, and thus compliance to the exercise program. NIVS delivery during exercise is labor intensive and may consequently increase the cost of the pulmonary rehabilitation program. NIVS should therefore only be considered in selected patients with COPD who demonstrate acute benefits from this intervention.

Heliox

The complex configuration of the bronchial tree, together with its branching angles and the internal airway diameter with its degree of roughness causes airflow to change from a turbulent to a laminar flow pattern as air moves from the central to the conductive and peripheral airways.^{148,149} Turbulent flow is further increased in patients with COPD consequent to airway inflammation and a loss of alveolar tethering, which causes narrowing of the airways. The resultant effect in such patients is an increased airway resistance and increased work of breathing at rest that becomes even more prominent during exercise.¹⁵⁰

Breathing a low-density gas mixture, such as normoxic heliox (He–O₂) – a mixture of 79% helium and 21% O₂ (79% He–21% O₂), decreases airway resistance by maintaining or re-establishing laminar flow within the tracheobronchial tree at higher flow rates.^{151–157} Similar to NIVS, heliox can be delivered non-invasively using different delivery

methods with a variety of patient interfaces such as a mouthpiece or facemask. The gas mixture is available in tanks of different sizes, with the 50L tank being the most frequently used. The tanks are pressurized at approximately 200 bar for a normoxic He–O₂ mixture, and the air regulators are connected to the ventilator¹⁵⁶ from which the mixture is delivered to the patient.

Effect of heliox on respiratory and peripheral muscles and their interaction

The administration of heliox, which is approximately three times less dense than air, during exercise in individuals with airflow obstruction has been shown to reduce resistive work of breathing and increase maximum expiratory flow, thus promoting faster lung emptying.^{40,158} In addition, evidence shows that heliox breathing increases V_E during exercise^{40,152,159–162} and also improves exercise tolerance,^{40,152,159,161,162} while reducing dynamic hyperinflation and dyspnea at isotime.^{41,152,159,160,162} This indicates that heliox is able to alleviate dyspnea and the work of breathing by primarily reducing ventilatory constraints.^{40,41,152,162} Using esophageal and gastric balloon catheters, Vogiatzis et al.¹⁶⁰ recently demonstrated the positive effect of heliox on reducing the work of the inspiratory and expiratory muscles during exercise.¹⁶⁰

There is emerging evidence which suggests that heliox-induced respiratory muscle unloading also improves distribution of the CO to the peripheral muscles during bicycle exercise in COPD patients.^{10,103,159,160} Richardson et al.¹⁰ showed that heliox administration promoted an increased VO_{2peak} and peak work load during whole body cycling exercise, with no change in arterial O₂ saturation, suggesting that the VO_2 increased secondary to enhanced peripheral O₂ availability and improved perfusion of the peripheral muscles.

Respiratory muscle unloading via heliox administration has also been associated with an improved O₂ delivery and extraction in the exercising locomotory muscles in moderate-to-severe COPD.⁴⁰ The increase in peripheral muscle O₂ delivery has been assumed to result from a redistribution of blood flow from the respiratory to the leg muscles.^{10,40} Interestingly, a recent study found that heliox breathing during near-maximum exercise (i.e., 75% peak work load) improved both quadriceps and intercostal muscle O₂ delivery due to an increase in both arterial O₂ content and quadriceps and intercostal muscle blood flow in patients with moderate-to-severe COPD with static but not dynamic hyperinflation.¹⁶⁰ In contrast to the previous studies, these findings do not support the “respiratory steal” phenomenon. Instead, it was concluded that the increased muscle blood flow and perfusion was due to a reduction in the work performed by the respiratory muscles.

In addition to the normoxic heliox, the effects of different O₂ concentrations (hyperoxia) in the heliox gas mixture have also been investigated. In these studies, improvement in exercise performance was associated with increased ventilatory capacity and decreased dynamic hyperinflation¹⁵² and dyspnea.^{41,152} These studies indicate that compared to either normoxic heliox or hyperoxia

alone, administration of a combination of helium and hyperoxia may provide a greater effect in reducing dynamic hyperinflation and work of breathing (WOB) and improving exercise performance.

Heliox breathing, similar to NIVS, unloads the respiratory muscles and relieves both dyspnea and leg discomfort during exercise. This allows COPD patients to exercise longer prior to exhaustion and enhances the physiological training effect,¹⁶² which in turn, could ultimately result in an improved activity of daily life and HRQoL.^{41,160} Despite the current findings, the overall cost of the ventilator set-up and the gas mixture makes the use of this therapy cumbersome/impractical and/or too expensive to be incorporated into routine pulmonary rehabilitation programs or training at home. Notwithstanding the evidence to support the use of heliox as an adjunct to exercise, further studies are needed to identify those individuals most likely to benefit from this intervention. Furthermore, studies are needed to determine the long-term utility of heliox during rehabilitation programs in COPD patients and determine how best to incorporate the latter into routine clinical practice.

Supplemental oxygen

In certain individuals with COPD, ventilation-perfusion mismatch and hypoventilation,¹⁶³ can lead to impaired gas exchange and hypoxemia¹⁶⁴ at rest and/or during exercise.^{112,165} Studies have shown that in patients with severe COPD, even routine daily activities such as walking, stair-climbing, washing, or eating can induce hypoxemia.^{166,167} Hypoxemia stimulates ventilatory drive, with the goal of increasing V_E , lowering PaCO₂ and in turn causing vasodilatation of the vascular bed, tachycardia, and an increased CO.^{112,168} Chronic hypoxemia can additionally lead to pulmonary hypertension and cor pulmonale (i.e. right heart failure), thereby reducing CO and impairing O₂ delivery.¹⁶⁹ With these factors compounding the effects of hyperinflation, it becomes evident that the hypoxemic patient with COPD is especially susceptible to lactic acidosis, muscle fatigue and reduce exercise capacity.^{112,169}

Effect of supplemental oxygen on the ventilatory and peripheral muscles and their interaction

Administration of supplemental O₂ during exercise to patients with COPD who are hypoxemic at rest and/or who desaturate with exercise has been shown to reduce V_E ¹⁷⁰, RR and ventilatory drive^{171–173} for a given exercise work load. The lower V_E which occurs secondary to a lower RR, has been reported to promote a reduction in dynamic hyperinflation,¹⁷² thus placing the diaphragm on a more optimal contractile portion of its length–tension curve. O₂ supplementation in hypoxemic patients has been reported to improve the diaphragm’s ability to sustain dynamic work,¹⁷⁴ to increase exercise endurance and to delay the onset of respiratory muscle fatigue¹⁷⁵ and dyspnea.¹⁷⁶ Interestingly, the association between increased endurance time with supplemental O₂ and the delayed onset of diaphragmatic fatigue has also been found in several studies that examined healthy individuals breathing against

an inspiratory resistance.^{177,178} While some authors have suggested that the lower V_E that occurs with supplemental O_2 is related to slower ventilatory kinetics in such hypoxemic patients,¹⁷¹ others have attributed the decreased V_E to delayed lactate accumulation, secondary to an increased peripheral muscle O_2 delivery, both in hypoxemic¹⁷⁹ and non-hypoxemic patients.¹⁸⁰ Evidence for the latter comes from a strong correlation between the decrease in V_E and fall in lactate accumulation ($r = 0.88$, $p = 0.001$).¹⁷⁹

Similar to the findings in hypoxemic patients, several studies have likewise reported reductions in the RR, V_E ,⁴² respiratory drive and dynamic hyperinflation^{42,43,180} along with improvements in exercise tolerance^{180,181} in normoxic COPD patients receiving supplemental O_2 during exercise. The mechanism linking reduced respiratory drive and improved exercise tolerance is said to be the prolongation of expiratory duration which reduces dynamic hyperinflation and the elastic work of breathing.^{42,43,180} In addition, it was shown that the decreased ventilation observed with supplemental O_2 during exercise in normoxic patients resulted in an increased mean femoral O_2 delivery, suggesting that a part of the blood flow may have been redistributed from the ventilatory to the peripheral muscles.¹⁸² However, the increased peripheral muscle blood flow during exercise may not necessarily be the result of a blood flow redistribution mechanism, since a concomitant increased inspiratory muscle blood flow has also been found with adjunct therapies that decrease the work of breathing, suggesting that other factors/mechanisms may be implicated.¹⁶⁰ Interestingly, Siqueira et al.¹⁸³ recently showed that despite improved central O_2 delivery and blood oxygenation with supplemental O_2 administration, some normoxic COPD patients do not benefit from O_2 supplementation during exercise due to an impaired intramuscular O_2 utilization.

Although the current evidence demonstrates that supplemental O_2 can improve O_2 saturation and peripheral tissue oxygenation, reduce dyspnea, and increase exercise capacity in both hypoxemic and non-hypoxemic COPD patients, the effects vary considerably among individual patients. Interestingly, Emtner et al.⁴² found that non-hypoxemic patients, who acutely improved exercise tolerance with supplemental O_2 , benefited more from using this therapy during exercise training in a pulmonary rehabilitation program. However, use of supplemental O_2 during exercise training for non-hypoxemic patients is not routine clinical practice at this point in time.

Conclusion

Although considerable research has been devoted to the effect of adjunct therapies for exercise training that may be useful in pulmonary rehabilitation programs, less is known about which patients are most likely to benefit from them. Reducing the work of breathing, dyspnea and peripheral muscle fatigue in patients with COPD is a key mechanism for improving exercise tolerance and activity. In the current review, three physiologically based interventions able to improve exercise tolerance have been discussed. It should be noted, however, that none of these therapies is currently routinely used in the context of

pulmonary rehabilitation, apart from supplemental O_2 for patients who have resting and/or exercise-induced O_2 desaturation, given that none has been proven to improve overall magnitude and/or duration of gains made in the context of routine clinical pulmonary rehabilitation programs. Although the administration of non-invasive ventilatory support, heliox and supplemental O_2 during exercise have been shown to unload the inspiratory muscles, reduce breathlessness, and enhance exercise endurance in patients with moderate-to-severe COPD, current available data demonstrates significant variability in their effectiveness across patients. Whether or not the symptoms limiting exercise contribute to such variability is unknown, raising the question whether patients who are limited by dyspnea obtain greater benefits from these adjunct therapies during exercise than those who are limited by leg fatigue. We propose that these techniques should be targeted towards individuals who show the most promising response. Examination of the acute effects of these adjunct therapies on exercise may provide insight into why some patients experience a greater benefit in use of these adjunct therapies during exercise training than others and may help to identify the most appropriate candidates for these forms of therapy.

Conflict of interest

None of the authors, A.M.M., M.d.M., D.S., and J.S., have any financial or other relationships that would constitute a conflict of interest.

References

1. Saetta M, Finkelstein R, Cosio MG. Morphological and cellular basis for airflow limitation in smokers. *Eur Respir J* 1994;7: 1505–15.
2. Global strategy for the diagnosis, management and prevention of COPD; 2009. Available from: <http://www.goldcopd.org> [updated January 16; cited 2011 January 16].
3. Becklake MR. Occupational exposures: evidence for a causal association with chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1989;140:S85–91.
4. Larson RK, Barman ML, Kueppers F, Fudenberg HH. Genetic and environmental determinants of chronic obstructive pulmonary disease. *Ann Intern Med* 1970;72:627–32.
5. Sandford AJ, Pare PD. Genetic risk factors for chronic obstructive pulmonary disease. *Clin Chest Med* 2000;21: 633–43.
6. Turato G, Zuin R, Saetta M. Pathogenesis and pathology of COPD. *Respiration* 2001;68:117–28.
7. Gallagher CG. Exercise limitation and clinical exercise testing in chronic obstructive pulmonary disease. *Clin Chest Med* 1994;15:305–26.
8. O'Donnell DE, Bertley JC, Chau LK, Webb KA. Qualitative aspects of exertional breathlessness in chronic airflow limitation: pathophysiologic mechanisms. *Am J Respir Crit Care Med* 1997;155:109–15.
9. Gosker HR, Lencer NH, Franssen FM, van der Vusse GJ, Wouters EF, Schols AM. Striking similarities in systemic factors contributing to decreased exercise capacity in patients with severe chronic heart failure or COPD. *Chest* 2003;123: 1416–24.

10. Richardson RS, Sheldon J, Poole DC, Hopkins SR, Ries AL, Wagner PD. Evidence of skeletal muscle metabolic reserve during whole body exercise in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1999;159:881–5.
11. Sala E, Roca J, Marrades RM, Alonso J, Gonzalez De Suso JM, Moreno A, et al. Effects of endurance training on skeletal muscle bioenergetics in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1999;159:1726–34.
12. Carter R, Holiday DB, Stocks J, Tjep B. Peak physiologic responses to arm and leg ergometry in male and female patients with airflow obstruction. *Chest* 2003;124:511–8.
13. Carlson DJ, Ries AL, Kaplan RM. Prediction of maximum exercise tolerance in patients with COPD. *Chest* 1991;100:307–11.
14. Gilbert R, Keighley J, Auchincloss Jr JH. Disability in patients with obstructive pulmonary disease. *Am Rev Respir Dis* 1964;90:383–94.
15. Jones NL, Jones G, Edwards RH. Exercise tolerance in chronic airway obstruction. *Am Rev Respir Dis* 1971;103:477–91.
16. Pepin V, Saey D, Laviolette L, Maltais F. Exercise capacity in chronic obstructive pulmonary disease: mechanisms of limitation. *COPD* 2007;4:195–204.
17. Killian KJ, LeBlanc P, Martin DH, Summers E, Jones NL, Campbell EJ. Exercise capacity and ventilatory, circulatory, and symptom limitation in patients with chronic airflow limitation. *Am Rev Respir Dis* 1992;146:935–40.
18. Laveneziana P, Wadell K, Webb K, O'Donnell DE. Exercise limitation in chronic obstructive pulmonary disease. *Curr Resp Med Rev* 2008;4:258–69.
19. Pepin V, Saey D, Whittom F, LeBlanc P, Maltais F. Walking versus cycling: sensitivity to bronchodilation in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2005;172:1517–22.
20. Deschenes D, Pepin V, Saey D, LeBlanc P, Maltais F. Locus of symptom limitation and exercise response to bronchodilation in chronic obstructive pulmonary disease. *J Cardiopulm Rehabil Prev* 2008;28:208–14.
21. Troosters T, Casaburi R, Gosselink R, Decramer M. Pulmonary rehabilitation in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2005;172:19–38.
22. Nici L, Donner C, Wouters E, Zuwallack R, Ambrosino N, Bourbeau J, et al. American Thoracic Society/European Respiratory Society statement on pulmonary rehabilitation. *Am J Respir Crit Care Med* 2006;173:1390–413.
23. Ries AL, Bauldoff GS, Carlin BW, Casaburi R, Emery CF, Mahler DA, et al. Pulmonary rehabilitation: joint ACCP/AACVPR evidence-based clinical practice guidelines. *Chest* 2007;131:45–42S.
24. O'Donnell DE, Aaron S, Bourbeau J, Hernandez P, Marciniuk DD, Balter M, et al. Canadian Thoracic Society recommendations for management of chronic obstructive pulmonary disease – 2007 update. *Can Respir J* 2007;14(Suppl. B):5B–32B.
25. Lacasse Y, Guyatt GH, Goldstein RS. The components of a respiratory rehabilitation program: a systematic overview. *Chest* 1997;111:1077–88.
26. American Thoracic Society. Pulmonary rehabilitation – 1999. *Am J Respir Crit Care Med* 1999;159:1666–82.
27. Lacasse Y, Goldstein R, Lasserion TJ, Martin S. Pulmonary rehabilitation for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2006;CD003793.
28. Bernard S, Whittom F, LeBlanc P, Jobin J, Belleau R, Berube C, et al. Aerobic and strength training in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1999;159:896–901.
29. Casaburi R, Patessio A, Ioli F, Zanaboni S, Donner CF, Wasserman K. Reductions in exercise lactic acidosis and ventilation as a result of exercise training in patients with obstructive lung disease. *Am Rev Respir Dis* 1991;143:9–18.
30. Casaburi R, Porszasz J, Burns MR, Carithers ER, Chang RS, Cooper CB. Physiologic benefits of exercise training in rehabilitation of patients with severe chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1997;155:1541–51.
31. Maltais F, LeBlanc P, Simard C, Jobin J, Berube C, Bruneau J, et al. Skeletal muscle adaptation to endurance training in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1996;154:442–7.
32. Vogiatzis I, Stratakis G, Simoes DC, Terzis G, Georgiadou O, Roussos C, et al. Effects of rehabilitative exercise on peripheral muscle TNF α , IL-6, IGF-I and MyoD expression in patients with COPD. *Thorax* 2007;62:950–6.
33. Takigawa N, Tada A, Soda R, Takahashi S, Kawata N, Shibayama T, et al. Comprehensive pulmonary rehabilitation according to severity of COPD. *Respir Med* 2007;101:326–32.
34. Coppoolse R, Schols AM, Baarends EM, Mostert R, Akkermans MA, Janssen PP, et al. Interval versus continuous training in patients with severe COPD: a randomized clinical trial. *Eur Respir J* 1999;14:258–63.
35. Cortopassi F, Castro AA, Porto EF, Colucci M, Fonseca G, Torre-Bouscoulet L, et al. Comprehensive exercise training improves ventilatory muscle function and reduces dyspnea perception in patients with COPD. *Monaldi Arch Chest Dis* 2009;71:106–12.
36. Gosselink R, Troosters T, Decramer M. Exercise training in COPD patients: the basic questions. *Eur Respir J* 1997;10:2884–91.
37. Maltais F, LeBlanc P, Jobin J, Berube C, Bruneau J, Carrier L, et al. Intensity of training and physiologic adaptation in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1997;155:555–61.
38. O'Donnell DE, Sanii R, Younes M. Improvement in exercise endurance in patients with chronic airflow limitation using continuous positive airway pressure. *Am Rev Respir Dis* 1988;138:1510–4.
39. Ambrosino N, Strambi S. New strategies to improve exercise tolerance in chronic obstructive pulmonary disease. *Eur Respir J* 2004;24:313–22.
40. Chiappa GR, Queiroga Jr F, Meda E, Ferreira LF, Diefenthaler F, Nunes M, et al. Heliox improves oxygen delivery and utilization during dynamic exercise in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2009;179:1004–10.
41. Laude EA, Duffy NC, Baveystock C, Dougill B, Campbell MJ, Lawson R, et al. The effect of helium and oxygen on exercise performance in chronic obstructive pulmonary disease: a randomized crossover trial. *Am J Respir Crit Care Med* 2006;173:865–70.
42. Emtner M, Porszasz J, Burns M, Somfay A, Casaburi R. Benefits of supplemental oxygen in exercise training in nonhypoxemic chronic obstructive pulmonary disease patients. *Am J Respir Crit Care Med* 2003;168:1034–42.
43. Somfay A, Porszasz J, Lee SM, Casaburi R. Dose-response effect of oxygen on hyperinflation and exercise endurance in nonhypoxaemic COPD patients. *Eur Respir J* 2001;18:77–84.
44. Rossi A, Polese G, Brandi G, Conti G. Intrinsic positive end-expiratory pressure (PEEPi). *Intensive Care Med* 1995;21:522–36.
45. Potter WA, Olafsson S, Hyatt RE. Ventilatory mechanics and expiratory flow limitation during exercise in patients with obstructive lung disease. *J Clin Invest* 1971;50:910–9.
46. Saetta M, Di Stefano A, Maestrelli P, Mapp CE, Ciaccia A, Fabbri LM. Structural aspects of airway inflammation in COPD. *Monaldi Arch Chest Dis* 1994;49:43–5.
47. Aliverti A, Iandelli I, Duranti R, Cala SJ, Kayser B, Kelly S, et al. Respiratory muscle dynamics and control during

- exercise with externally imposed expiratory flow limitation. *J Appl Physiol* 2002;**92**:1953–63.
48. O'Donnell DE, Revill SM, Webb KA. Dynamic hyperinflation and exercise intolerance in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2001;**164**:770–7.
 49. Pride NB, Macklem PT. Lung mechanics in disease. In: Macklem PT, Mead J, editors. *Handbook of physiology, section 3: the respiratory system*. Bethesda: American Physiological Society; 1986. p. 659–92.
 50. O'Donnell DE. Hyperinflation, dyspnea, and exercise intolerance in chronic obstructive pulmonary disease. *Proc Am Thorac Soc*. 2006;**3**:180–4.
 51. Rochester DF, Braun NM. Determinants of maximal inspiratory pressure in chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1985;**132**:42–7.
 52. O'Donnell DE, Laveneziana P. Dyspnea and activity limitation in COPD: mechanical factors. *COPD* 2007;**4**:225–36.
 53. Loring SH, Garcia-Jacques M, Malhotra A. Pulmonary characteristics in COPD and mechanisms of increased work of breathing. *J Appl Physiol* 2009;**107**:309–14.
 54. De Troyer A, Wilson TA. Effect of acute inflation on the mechanics of the inspiratory muscles. *J Appl Physiol* 2009;**107**:315–23.
 55. Sinderby C, Beck J, Spahija J, Weinberg J, Grassino A. Voluntary activation of the human diaphragm in health and disease. *J Appl Physiol* 1998;**85**:2146–58.
 56. Sinderby C, Spahija J, Beck J, Kaminski D, Yan S, Comtois N, et al. Diaphragm activation during exercise in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2001;**163**:1637–41.
 57. Spahija J, Beck J, Lindstrom L, Begin P, de Marchie M, Sinderby C. Effect of increased diaphragm activation on diaphragm power spectrum center frequency. *Respir Physiol Neurobiol*. 2005;**146**:67–76.
 58. Spahija J, Beck J, Sinderby C. Respiratory failure and diaphragm function. In: Vicente JL, editor. *Yearbook of intensive care and emergency medicine*. Berlin, Heidelberg, New York: Springer-Verlag; 2004. p. 325–32.
 59. Shindoh C, Hida W, Kikuchi Y, Taguchi O, Miki H, Takishima T, et al. Oxygen consumption of respiratory muscles in patients with COPD. *Chest* 1994;**105**:790–7.
 60. Sinderby C, Spahija J, Beck J. Changes in respiratory effort sensation over time are linked to the frequency content of diaphragm electrical activity. *Am J Respir Crit Care Med* 2001;**163**:905–10.
 61. Levison H, Cherniack RM. Ventilatory cost of exercise in chronic obstructive pulmonary disease. *J Appl Physiol* 1968;**25**:21–7.
 62. Gosker HR, Wouters EF, van der Vusse GJ, Schols AM. Skeletal muscle dysfunction in chronic obstructive pulmonary disease and chronic heart failure: underlying mechanisms and therapy perspectives. *Am J Clin Nutr* 2000;**71**:1033–47.
 63. Casaburi R. Deconditioning. In: Fishman A, editor. *Pulmonary rehabilitation*. New York: Marcel Dekker; 1996. p. 213–30.
 64. Schols AM, Buurman WA, Staal van den Brekel AJ, Dentener MA, Wouters EF. Evidence for a relation between metabolic derangements and increased levels of inflammatory mediators in a subgroup of patients with chronic obstructive pulmonary disease. *Thorax* 1996;**51**:819–24.
 65. Saltin B, Gollnick P. In: Peachey L, editor. *Skeletal muscle adaptability: significance for metabolism and performance*. Washington, DC: American Physiological Society; 1986.
 66. Whittom F, Jobin J, Simard PM, Leblanc P, Simard C, Bernard S, et al. Histochemical and morphological characteristics of the vastus lateralis muscle in patients with chronic obstructive pulmonary disease. *Med Sci Sports Exerc* 1998;**30**:1467–74.
 67. Jakobsson P, Jorfeldt L, Henriksson J. Metabolic enzyme activity in the quadriceps femoris muscle in patients with severe chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1995;**151**:374–7.
 68. Bernard S, LeBlanc P, Whittom F, Carrier G, Jobin J, Belleau R, et al. Peripheral muscle weakness in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1998;**158**:629–34.
 69. Hildebrand IL, Sylven C, Esbjornsson M, Hellstrom K, Jansson E. Does chronic hypoxaemia induce transformations of fibre types? *Acta Physiol Scand* 1991;**141**:435–9.
 70. Maltais F, LeBlanc P, Whittom F, Simard C, Marquis K, Belanger M, et al. Oxidative enzyme activities of the vastus lateralis muscle and the functional status in patients with COPD. *Thorax* 2000;**55**:848–53.
 71. Gosselink R, Troosters T, Decramer M. Peripheral muscle weakness contributes to exercise limitation in COPD. *Am J Respir Crit Care Med* 1996;**153**:976–80.
 72. Edwards RH, Young A, Hosking GP, Jones DA. Human skeletal muscle function: description of tests and normal values. *Clin Sci Mol Med* 1977;**52**:283–90.
 73. Edwards RH. Human muscle function and fatigue. In: Porter H, Whelan J, editors. *Human muscle fatigue: physiological mechanisms*. London: Pitman Medical; 1981. p. 1–18.
 74. Polkey MI, Kyroussis D, Hammegard CH, Mills GH, Green M, Moxham J. Quadriceps strength and fatigue assessed by magnetic stimulation of the femoral nerve in man. *Muscle Nerve* 1996;**19**:549–55.
 75. Jeffery Mador M, Kufel TJ, Pineda L. Quadriceps fatigue after cycle exercise in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2000;**161**:447–53.
 76. Mador MJ, Kufel TJ, Pineda LA, Steinwald A, Aggarwal A, Upadhyay AM, et al. Effect of pulmonary rehabilitation on quadriceps fatigability during exercise. *Am J Respir Crit Care Med* 2001;**163**:930–5.
 77. Saey D, Debigare R, LeBlanc P, Mador MJ, Cote CH, Jobin J, et al. Contractile leg fatigue after cycle exercise: a factor limiting exercise in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2003;**168**:425–30.
 78. Man WD, Soliman MG, Gearing J, Radford SG, Rafferty GF, Gray BJ, et al. Symptoms and quadriceps fatigability after walking and cycling in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2003;**168**:562–7.
 79. Hamilton AL, Killian KJ, Summers E, Jones NL. Muscle strength, symptom intensity, and exercise capacity in patients with cardiorespiratory disorders. *Am J Respir Crit Care Med* 1995;**152**:2021–31.
 80. Casaburi R. Exercise training in chronic obstructive lung disease. In: Casaburi R, Petty TL, editors. *Principles and practice of pulmonary rehabilitation*. Philadelphia: Saunders; 1993. p. 202–24.
 81. Maltais F, Jobin J, Sullivan MJ, Bernard S, Whittom F, Killian KJ, et al. Metabolic and hemodynamic responses of lower limb during exercise in patients with COPD. *J Appl Physiol* 1998;**84**:1573–80.
 82. Weisman M, Zeballos R. Clinical exercise testing. In: Bolliger C, editor. *Prog respir res*. Basel: Karger; 2002. p. 138–58.
 83. Guyton AC, Hall JE. Heart muscle; the heart as a pump. In: Guyton AC, Hall JE, editors. *Textbook of medical physiology*. 11th ed. Philadelphia: Elsevier Saunders Inc.; 2006. p. 99.
 84. Bassett Jr DR, Howley ET. Limiting factors for maximum oxygen uptake and determinants of endurance performance. *Med Sci Sports Exerc* 2000;**32**:70–84.
 85. Faulkner JA, Heigenhauser GJ, Schork MA. The cardiac output–oxygen uptake relationship of men during graded bicycle ergometry. *Med Sci Sports Exerc* 1977;**9**:148–54.
 86. Proctor DN, Beck KC, Shen PH, Eickhoff TJ, Halliwill JR, Joyner MJ. Influence of age and gender on cardiac

- output- VO_2 relationships during submaximal cycle ergometry. *J Appl Physiol* 1998;**84**:599–605.
87. Stewart RI, Lewis CM. Cardiac output during exercise in patients with COPD. *Chest* 1986;**89**:199–205.
 88. Bogaard HJ, Dekker BM, Arntzen BW, Woltjer HH, van Keimpema AR, Postmus PE, et al. The haemodynamic response to exercise in chronic obstructive pulmonary disease: assessment by impedance cardiography. *Eur Respir J* 1998;**12**:374–9.
 89. Morrison DA, Adcock K, Collins CM, Goldman S, Caldwell JH, Schwarz MI. Right ventricular dysfunction and the exercise limitation of chronic obstructive pulmonary disease. *J Am Coll Cardiol* 1987;**9**:1219–29.
 90. Light RW, Mintz HM, Linden GS, Brown SE. Hemodynamics of patients with severe chronic obstructive pulmonary disease during progressive upright exercise. *Am Rev Respir Dis* 1984;**130**:391–5.
 91. Sietsema K. Cardiovascular limitations in chronic pulmonary disease. *Med Sci Sports Exerc* 2001;**33**:S656–61.
 92. Jorgensen K, Muller MF, Nel J, Upton RN, Houtz E, Ricksten SE. Reduced intrathoracic blood volume and left and right ventricular dimensions in patients with severe emphysema: an MRI study. *Chest* 2007;**131**:1050–7.
 93. Holverda S, Rietema H, Westerhof N, Marcus JT, Gan CT, Postmus PE, et al. Stroke volume increase to exercise in chronic obstructive pulmonary disease is limited by increased pulmonary artery pressure. *Heart* 2009;**95**:137–41.
 94. Matthay RA, Arroliga AC, Wiedemann HP, Schulman DS, Mahler DA. Right ventricular function at rest and during exercise in chronic obstructive pulmonary disease. *Chest* 1992;**101**:2555–62S.
 95. Baril J, de Souza M, Leroy D, Ofir D, Aguilaniu B, Gladys C, et al. Does dynamic hyperinflation impair submaximal exercise cardiac output in chronic obstructive pulmonary disease? *Clin Invest Med* 2006;**29**:104–9.
 96. Sabapathy S, Awater MF, Schneider DA, Kingsley RA, Hopman MT, Morris NR. Lower limb vasodilatory capacity is not reduced in patients with moderate COPD. *Int J Chron Obstruct Pulmon Dis* 2006;**1**:73–81.
 97. Oelberg DA, Medoff BD, Markowitz DH, Pappagianopoulos PP, Ginns LC, Systrom DM. Systemic oxygen extraction during incremental exercise in patients with severe chronic obstructive pulmonary disease. *Eur J Appl Physiol Occup Physiol* 1998;**78**:201–7.
 98. Vassaux C, Torre-Bouscoulet L, Zeineldine S, Cortopassi F, Paz-Diaz H, Celli BR, et al. Effects of hyperinflation on the oxygen pulse as a marker of cardiac performance in COPD. *Eur Respir J* 2008;**32**:1275–82.
 99. Montes de Oca M, Rassulo J, Celli BR. Respiratory muscle and cardiopulmonary function during exercise in very severe COPD. *Am J Respir Crit Care Med* 1996;**154**:1284–9.
 100. Tutar E, Kaya A, Gulec S, Ertas F, Erol C, Ozdemir O, et al. Echocardiographic evaluation of left ventricular diastolic function in chronic cor pulmonale. *Am J Cardiol* 1999;**83**:1414–7. A9.
 101. Barr RG, Bluemke DA, Ahmed FS, Carr JJ, Enright PL, Hoffman EA, et al. Percent emphysema, airflow obstruction, and impaired left ventricular filling. *N Engl J Med* 2010;**362**:217–27.
 102. Watz H, Waschki B, Meyer T, Kretschmar G, Kirsten A, Claussen M, et al. Decreasing cardiac chamber sizes and associated heart dysfunction in COPD: role of hyperinflation. *Chest* 2010;**138**:32–8.
 103. Borghi-Silva A, Oliveira CC, Carrascosa C, Maia J, Berton DC, Queiroga Jr F, et al. Respiratory muscle unloading improves leg muscle oxygenation during exercise in patients with COPD. *Thorax* 2008;**63**:910–5.
 104. Harms CA, Babcock MA, McClaran SR, Pegelow DF, Nickele GA, Nelson WB, et al. Respiratory muscle work compromises leg blood flow during maximal exercise. *J Appl Physiol* 1997;**82**:1573–83.
 105. Harms CA, Wetter TJ, St Croix CM, Pegelow DF, Dempsey JA. Effects of respiratory muscle work on exercise performance. *J Appl Physiol* 2000;**89**:131–8.
 106. Simon M, LeBlanc P, Jobin J, Desmeules M, Sullivan MJ, Maltais F. Limitation of lower limb VO_2 during cycling exercise in COPD patients. *J Appl Physiol* 2001;**90**:1013–9.
 107. Stark-Leyva KN, Beck KC, Johnson BD. Influence of expiratory loading and hyperinflation on cardiac output during exercise. *J Appl Physiol* 2004;**96**:1920–7.
 108. Peter JV, Moran JL, Phillips-Hughes J, Warn D. Noninvasive ventilation in acute respiratory failure – a meta-analysis update. *Crit Care Med* 2002;**30**:555–62.
 109. Pevernagie D, Janssens JP, De Backer W, Elliott M, Pepperell J, Andreas S. Ventilatory support and pharmacological treatment of patients with central apnoea or hypoventilation during sleep. *Eur Respir Rev* 2007;**16**:115–24.
 110. Barakat S, Michele G, Nesme P, Nicole V, Guy A. Effect of a noninvasive ventilatory support during exercise of a program in pulmonary rehabilitation in patients with COPD. *Int J Chron Obstruct Pulmon Dis* 2007;**2**:585–91.
 111. Schiller O, Schonfeld T, Yaniv I, Stein J, Kadmon G, Nahum E. Bi-level positive airway pressure ventilation in pediatric oncology patients with acute respiratory failure. *J Intensive Care Med* 2009;**24**:383–8.
 112. Hoo GW. Nonpharmacologic adjuncts to training during pulmonary rehabilitation: the role of supplemental oxygen and noninvasive ventilation. *J Rehabil Res Dev*. 2003;**40**:81–97.
 113. Ambrosino N, Palmiero G, Strambi SK. New approaches in pulmonary rehabilitation. *Clin Chest Med* 2007;**28**:629–38. vii.
 114. O'Donnell DE, Sanii R, Giesbrecht G, Younes M. Effect of continuous positive airway pressure on respiratory sensation in patients with chronic obstructive pulmonary disease during submaximal exercise. *Am Rev Respir Dis* 1988;**138**:1185–91.
 115. Costa D, Toledo A, Silva AB, Sampaio LM. Influence of noninvasive ventilation by BiPAP on exercise tolerance and respiratory muscle strength in chronic obstructive pulmonary disease patients (COPD). *Rev Lat Am Enfermagem* 2006;**14**:378–82.
 116. Borghi-Silva A, Reis MS, Mendes RG, Pantoni CB, Simoes RP, Martins LE, et al. Noninvasive ventilation acutely modifies heart rate variability in chronic obstructive pulmonary disease patients. *Respir Med* 2008;**102**:1117–23.
 117. Borghi-Silva A, Di Thommazo L, Pantoni CB, Mendes RG, Salvini Tde F, Costa D. Non-invasive ventilation improves peripheral oxygen saturation and reduces fatigability of quadriceps in patients with COPD. *Respirology* 2009;**14**:537–44.
 118. Borghi-Silva A, Mendes RG, Toledo AC, Malosa Sampaio LM, da Silva TP, Kunikushita LN, et al. Adjuncts to physical training of patients with severe COPD: oxygen or noninvasive ventilation? *Respir Care* 2010;**55**:885–94.
 119. Toledo A, Borghi-Silva A, Sampaio LM, Ribeiro KP, Baldissera V, Costa D. The impact of noninvasive ventilation during the physical training in patients with moderate-to-severe chronic obstructive pulmonary disease (COPD). *Clinics (Sao Paulo)* 2007;**62**:113–20.
 120. Reuveny R, Ben-Dov I, Gaides M, Reichert N. Ventilatory support during training improves training benefit in severe chronic airway obstruction. *Isr Med Assoc J* 2005;**7**:151–5.
 121. Johnson JE, Gavin DJ, Adams-Dramiga S. Effects of training with heliox and noninvasive positive pressure ventilation on

- exercise ability in patients with severe COPD. *Chest* 2002; **122**:464–72.
122. Kohnlein T, Schonheit-Kenn U, Winterkamp S, Welte T, Kenn K. Noninvasive ventilation in pulmonary rehabilitation of COPD patients. *Respir Med* 2009; **103**:1329–36.
 123. Brochard L. Pressure support ventilation. In: Tobin MJ, editor. *Principles and practice of mechanical ventilation*. New York: McGraw-Hill Inc.; 1994. p. 239–56.
 124. van 't Hul A, Gosselink R, Hollander P, Postmus P, Kwakkel G. Training with inspiratory pressure support in patients with severe COPD. *Eur Respir J* 2006; **27**:65–72.
 125. van 't Hul A, Gosselink R, Hollander P, Postmus P, Kwakkel G. Acute effects of inspiratory pressure support during exercise in patients with COPD. *Eur Respir J* 2004; **23**:34–40.
 126. Keilty SE, Ponte J, Fleming TA, Moxham J. Effect of inspiratory pressure support on exercise tolerance and breathlessness in patients with severe stable chronic obstructive pulmonary disease. *Thorax* 1994; **49**:990–4.
 127. Maltais F, Reissmann H, Gottfried SB. Pressure support reduces inspiratory effort and dyspnea during exercise in chronic airflow obstruction. *Am J Respir Crit Care Med* 1995; **151**:1027–33.
 128. Younes M. Proportional assist ventilation, a new approach to ventilatory support. Theory. *Am Rev Respir Dis* 1992; **145**:114–20.
 129. Hernandez P, Maltais F, Gursahaney A, Leblanc P, Gottfried SB. Proportional assist ventilation may improve exercise performance in severe chronic obstructive pulmonary disease. *J Cardiopulm Rehabil* 2001; **21**:135–42.
 130. Dolmage TE, Goldstein RS. Proportional assist ventilation and exercise tolerance in subjects with COPD. *Chest* 1997; **111**:948–54.
 131. Hawkins P, Johnson LC, Nikolettou D, Hamnegard CH, Sherwood R, Polkey MI, et al. Proportional assist ventilation as an aid to exercise training in severe chronic obstructive pulmonary disease. *Thorax* 2002; **57**:853–9.
 132. Bianchi L, Foglio K, Porta R, Baiardi R, Vitacca M, Ambrosino N. Lack of additional effect of adjunct of assisted ventilation to pulmonary rehabilitation in mild COPD patients. *Respir Med* 2002; **96**:359–67.
 133. Spahija J, Albert M, Bellemare P, Lacoste G, De Marchie M. Downregulation of diaphragm activation with pressure support ventilation (PSV) in COPD and healthy subjects during exercise. *Am J Respir Crit Care Med* 2007; **175**:A224.
 134. Diaz O, Iglesia R, Ferrer M, Zavala E, Santos C, Wagner PD, et al. Effects of noninvasive ventilation on pulmonary gas exchange and hemodynamics during acute hypercapnic exacerbations of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1997; **156**:1840–5.
 135. Petrof BJ, Calderini E, Gottfried SB. Effect of CPAP on respiratory effort and dyspnea during exercise in severe COPD. *J Appl Physiol* 1990; **69**:179–88.
 136. Kyroussis D, Polkey MI, Hamnegard CH, Mills GH, Green M, Moxham J. Respiratory muscle activity in patients with COPD walking to exhaustion with and without pressure support. *Eur Respir J* 2000; **15**:649–55.
 137. Gorman RB, McKenzie DK, Butler JE, Tolman JF, Gandevia SC. Diaphragm length and neural drive after lung volume reduction surgery. *Am J Respir Crit Care Med* 2005; **172**:1259–66.
 138. Oliveira CC, Carrasosa CR, Borghi-Silva A, Berton DC, Queiroga Jr F, Ferreira EM, et al. Influence of respiratory pressure support on hemodynamics and exercise tolerance in patients with COPD. *Eur J Appl Physiol* 2010; **109**:681–9.
 139. Polkey MI, Hawkins P, Kyroussis D, Ellum SG, Sherwood R, Moxham J. Inspiratory pressure support prolongs exercise induced lactataemia in severe COPD. *Thorax* 2000; **55**:547–9.
 140. Bianchi L, Foglio K, Pagani M, Vitacca M, Rossi A, Ambrosino N. Effects of proportional assist ventilation on exercise tolerance in COPD patients with chronic hypercapnia. *Eur Respir J* 1998; **11**:422–7.
 141. Moga AM, de Marchie M, Delisle S, Spahija J. The effect of BiPAP on symptom-limited maximum exercise in COPD patients. *Eur Respir J* 2011; **38**:4809.
 142. Evison H, Cherniack RM. Ventilatory cost of exercise in chronic obstructive pulmonary disease. *J Appl Physiol* 1968; **25**:21–7.
 143. Harms CA, Wetter TJ, McClaran SR, Pegelow DF, Nিকেle GA, Nelson WB, et al. Effects of respiratory muscle work on cardiac output and its distribution during maximal exercise. *J Appl Physiol* 1998; **85**:609–18.
 144. Pinsky MR, Matuschak GM, Klain M. Determinants of cardiac augmentation by elevations in intrathoracic pressure. *J Appl Physiol* 1985; **58**:1189–98.
 145. Duke GJ, Bersten AD. Non-invasive ventilation for adult acute respiratory failure. Part I. *Crit Care Resusc* 1999; **1**:198.
 146. Kong W, Wang C, Yang Y, Huang K, Jiang C. Effects of extrinsic positive end-expiratory pressure on cardiopulmonary function in patients with chronic obstructive pulmonary disease. *Chin Med J (Engl)* 2001; **114**:912–5.
 147. Younes M, Puddy A, Roberts D, Light RB, Quesada A, Taylor K, et al. Proportional assist ventilation. Results of an initial clinical trial. *Am Rev Respir Dis* 1992; **145**:121–9.
 148. Papamoschou D. Theoretical validation of the respiratory benefits of helium–oxygen mixtures. *Respir Physiol* 1995; **99**:183–90.
 149. Yanai M, Sekizawa K, Ohru T, Sasaki H, Takishima T. Site of airway obstruction in pulmonary disease: direct measurement of intrabronchial pressure. *J Appl Physiol* 1992; **72**:1016–23.
 150. O'Donnell DE, Banzett RB, Carrieri-Kohlman V, Casaburi R, Davenport PW, Gandevia SC, et al. Pathophysiology of dyspnea in chronic obstructive pulmonary disease: a roundtable. *Proc Am Thorac Soc*. 2007; **4**:145–68.
 151. Martin D, Day J, Ward G, Carter E, Chesrown S. Effects of breathing a normoxic helium mixture on exercise tolerance of patients with cystic fibrosis. *Pediatr Pulmonol* 1994; **18**:206–10.
 152. Eves ND, Petersen SR, Haykowsky MJ, Wong EY, Jones RL. Helium-hyperoxia, exercise, and respiratory mechanics in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2006; **174**:763–71.
 153. Esposito F, Ferretti G. The effects of breathing He–O₂ mixtures on maximal oxygen consumption in normoxic and hypoxic men. *J Physiol* 1997; **503**(Pt 1):215–22.
 154. Hunt T, Williams MT, Frith P, Schembri D. Heliox, dyspnoea and exercise in COPD. *Eur Respir Rev* 2010; **19**:30–8.
 155. Eves N, Ford G. Helium–oxygen: a versatile therapy to “lighten the load” of chronic obstructive pulmonary disease (COPD). *Respir Med COPD Update* 2007; **3**:87–94.
 156. Gainnier M, Forel JM. Clinical review: use of helium–oxygen in critically ill patients. *Crit Care* 2006; **10**:241.
 157. Valli G, Paoletti P, Savi D, Martolini D, Palange P. Clinical use of Heliox in asthma and COPD. *Monaldi Arch Chest Dis* 2007; **67**:159–64.
 158. Babb TG. Breathing He–O₂ increases ventilation but does not decrease the work of breathing during exercise. *Am J Respir Crit Care Med* 2001; **163**:1128–34.
 159. Amann M, Regan MS, Kobitany M, Eldridge MW, Boutellier U, Pegelow DF, et al. Impact of pulmonary system limitations on locomotor muscle fatigue in patients with COPD. *Am J Physiol Regul Integr Comp Physiol* 2010; **299**:R314–24.
 160. Vogiatzis I, Habazettl H, Aliverti A, Athanasopoulos D, Louvaris Z, Lomauro A, et al. Effect of helium breathing on intercostal and quadriceps muscle blood flow during exercise in COPD patients. *Am J Physiol Regul Integr Comp Physiol* 2011; **300**:R1549–59.

161. Oelberg DA, Kacmarek RM, Pappagianopoulos PP, Ginns LC, Systrom DM. Ventilatory and cardiovascular responses to inspired He—O₂ during exercise in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1998;**158**:1876–82.
162. Palange P, Valli G, Onorati P, Antonucci R, Paoletti P, Rosato A, et al. Effect of heliox on lung dynamic hyperinflation, dyspnea, and exercise endurance capacity in COPD patients. *J Appl Physiol* 2004;**97**:1637–42.
163. Begin P, Grassino A. Inspiratory muscle dysfunction and chronic hypercapnia in chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1991;**143**:905–12.
164. Nonoyama ML, Brooks D, Lacasse Y, Guyatt GH, Goldstein RS. Oxygen therapy during exercise training in chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2007:CD005372.
165. Nonoyama ML, Brooks D, Guyatt GH, Goldstein RS. Effect of oxygen on health quality of life in patients with chronic obstructive pulmonary disease with transient exertional hypoxemia. *Am J Respir Crit Care Med* 2007;**176**:343–9.
166. Soguel Schenkel N, Burdet L, de Muralt B, Fitting JW. Oxygen saturation during daily activities in chronic obstructive pulmonary disease. *Eur Respir J* 1996;**9**:2584–9.
167. Dreher M, Waltersbacher S, Sonntag F, Prettin S, Kabitz HJ, Windisch W. Exercise in severe COPD: is walking different from stair-climbing? *Respir Med* 2008;**102**:912–8.
168. Tarpy SP, Celli BR. Long-term oxygen therapy. *N Engl J Med* 1995;**333**:710–4.
169. O'Donohue W. Oxygen therapy in pulmonary rehabilitation. In: Hodgkin J, Celli B, Connors G, editors. *Pulmonary rehabilitation: guidelines for success*. 3rd ed. Philadelphia: Lippincott Williams & Wilkins; 2000. p. 135–46.
170. Mannix ET, Manfredi F, Palange P, Dowdeswell IR, Farber MO. Oxygen may lower the O₂ cost of ventilation in chronic obstructive lung disease. *Chest* 1992;**101**:910–5.
171. Payen JF, Wuyam B, Levy P, Reutenauer H, Stieglitz P, Paramelle B, et al. Muscular metabolism during oxygen supplementation in patients with chronic hypoxemia. *Am Rev Respir Dis* 1993;**147**:592–8.
172. O'Donnell DE, D'Arsigny C, Webb KA. Effects of hyperoxia on ventilatory limitation during exercise in advanced chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2001;**163**:892–8.
173. Light RW, Mahutte CK, Stansbury DW, Fischer CE, Brown SE. Relationship between improvement in exercise performance with supplemental oxygen and hypoxic ventilatory drive in patients with chronic airflow obstruction. *Chest* 1989;**95**:751–6.
174. Criner GJ, Celli BR. Ventilatory muscle recruitment in exercise with O₂ in obstructed patients with mild hypoxemia. *J Appl Physiol* 1987;**63**:195–200.
175. Bye PT, Esau SA, Levy RD, Shiner RJ, Macklem PT, Martin JG, et al. Ventilatory muscle function during exercise in air and oxygen in patients with chronic air-flow limitation. *Am Rev Respir Dis* 1985;**132**:236–40.
176. Dean NC, Brown JK, Himelman RB, Doherty JJ, Gold WM, Stulbarg MS. Oxygen may improve dyspnea and endurance in patients with chronic obstructive pulmonary disease and only mild hypoxemia. *Am Rev Respir Dis* 1992;**146**:941–5.
177. Bye PT, Esau SA, Walley KR, Macklem PT, Pardy RL. Ventilatory muscles during exercise in air and oxygen in normal men. *J Appl Physiol* 1984;**56**:464–71.
178. Pardy RL, Bye PT. Diaphragmatic fatigue in normoxia and hyperoxia. *J Appl Physiol* 1985;**58**:738–42.
179. O'Donnell DE, Bain DJ, Webb KA. Factors contributing to relief of exertional breathlessness during hyperoxia in chronic airflow limitation. *Am J Respir Crit Care Med* 1997;**155**:530–5.
180. Somfay A, Porszasz J, Lee SM, Casaburi R. Effect of hyperoxia on gas exchange and lactate kinetics following exercise onset in nonhypoxemic COPD patients. *Chest* 2002;**121**:393–400.
181. Voduc N, Tessier C, Sabri E, Fergusson D, Lavallee L, Aaron SD. Effects of oxygen on exercise duration in chronic obstructive pulmonary disease patients before and after pulmonary rehabilitation. *Can Respir J* 2010;**17**:e14–9.
182. Maltais F, Simon M, Jobin J, Desmeules M, Sullivan MJ, Belanger M, et al. Effects of oxygen on lower limb blood flow and O₂ uptake during exercise in COPD. *Med Sci Sports Exerc* 2001;**33**:916–22.
183. Siqueira AC, Borghi-Silva A, Bravo DM, Ferreira EM, Chiappa GR, Neder JA. Effects of hyperoxia on the dynamics of skeletal muscle oxygenation at the onset of heavy-intensity exercise in patients with COPD. *Respir Physiol Neurobiol*. 2010;**172**:8–14.